Unique Plan Description: DAPTOmycin (Cubicin) Adult
Plan Selection Display: DAPTOmycin (Cubicin) Adult
PlanType: Medical
Version: 1
Begin Effective Date:
End Effective Date: Current
Available at: ***BHMH***
C A DEAN HOSPIT
EMMC
MERCY HOSPITAL
NL MAINE COAST HOSPITAL
SVH
TAMC - THE AROO
Plan Comment:

DAPTOmycin (Cubicin) Adult
Medications

For use in patient 18 years old or greater.(NOTE)*
DAPTOmycin dose is based on adjusted body weight.(NOTE)*

Adjusted body weight calculator link - scroll down within the link to find the Adjusted Body Weight calculator

Adjusted Body Weight (kg) = 0.4 (Total Body Weight - Adjusted Body Weight) + Adjusted Body Weight(NOTE)*

Creatinine clearance can be found in Related Results.(NOTE)*
Skin & Soft Tissue Infections(NOTE)*
Adjusted body weight less than 40 kg(NOTE)*
CrCl 30 mL/min or greater(NOTE)*

☐ DAPTOmycin
300 mg, Soln, IV Push, Every 24 Hour Interval, CrCl 30 mL/min or greater
Comments: Adjusted body weight less than 40 kg and creatinine clearance 30 mL/min or greater

A pharmacy consult is recommended for adjusted body weight less than 40 kg(NOTE)*

☐ Antimicrobial - Pharmacy Consult
ONCE
Comments: A pharmacy consult is recommended for adjusted body weight less than 40 kg. Consult to be performed by infectious diseases pharmacist.

CrCl less than 30 mL/min(NOTE)*

☐ DAPTOmycin
300 mg, Soln, IV Push, Every 48 Hour Interval, CrCl less than 30 mL/min
Comments: Adjusted body weight less than 40 kg and creatinine clearance less than 30 mL/min

A pharmacy consult is recommended for adjusted body weight less than 40 kg(NOTE)*

☐ Antimicrobial - Pharmacy Consult
ONCE
Comments: A pharmacy consult is recommended for adjusted body weight less than 40 kg. Consult to be performed by infectious diseases pharmacist.

Adjusted body weight 40 up to 69 kg(NOTE)*
CrCl 30 mL/min or greater(NOTE)*

☐ DAPTOmycin
300 mg, Soln, IV Push, Every 24 Hour Interval, CrCl 30 mL/min or greater
Comments: Adjusted body weight 40 up to 69 kg and creatinine clearance 30 mL/min or greater

CrCl less than 30 mL/min(NOTE)*

☐ DAPTOmycin
300 mg, Soln, IV Push, Every 48 Hour Interval, CrCl less than 30 mL/min
Comments: Adjusted body weight 40 up to 69 kg and creatinine clearance less than 30 mL/min

Adjusted body weight 70 up to 100 kg(NOTE)*
CrCl 30 mL/min or greater(NOTE)*

☐ DAPTOmycin
500 mg, Soln, IV Push, Every 24 Hour Interval, CrCl 30 mL/min or greater
Comments: Adjusted body weight 70 up to 100 kg and creatinine clearance 30 mL/min or greater

CrCl less than 30 mL/min(NOTE)*

☐ DAPTOmycin
500 mg, Soln, IV Push, Every 48 Hour Interval, CrCl less than 30 mL/min
Comments: Adjusted body weight 70 up to 100 kg and creatinine clearance less than 30 mL/min

Adjusted body weight 101 kg or greater(NOTE)*
CrCl 30 mL/min or greater(NOTE)*

☐ DAPTOmycin
500 mg, Soln, IV Push, Every 24 Hour Interval, CrCl 30 mL/min or greater
Comments: Adjusted body weight 101 kg or greater and creatinine clearance 30 mL/min or greater

A pharmacy consult is recommended for adjusted body weight 101 kg or greater(NOTE)*

☐ Antimicrobial - Pharmacy Consult
ONCE
Comments: A pharmacy consult is recommended for adjusted body weight 101 kg or greater. Consult to be performed by infectious diseases pharmacist.

CrCl less than 30 mL/min(NOTE)*

☐ DAPTOmycin
500 mg, Soln, IV Push, Every 48 Hour Interval, CrCl less than 30 mL/min
Comments: Adjusted body weight 101 kg or greater and creatinine clearance less than 30 mL/min

A pharmacy consult is recommended for adjusted body weight 101 kg or greater(NOTE)*

☐ Antimicrobial - Pharmacy Consult
ONCE
Comments: A pharmacy consult is recommended for adjusted body weight 101 kg or greater. Consult to be performed by infectious diseases pharmacist.

Invasive MRSA Enterococcus faecalis & Bone and Joint Infections(NOTE)*
Adjusted body weight less than 40 kg(NOTE)*
CrCl 30 mL/min or greater(NOTE)*

☐ DAPTOmycin
300 mg, Soln, IV Push, Every 24 Hour Interval, CrCl 30 mL/min or greater
Comments: Adjusted body weight less than 40 kg and creatinine clearance 30 mL/min or greater

A pharmacy consult is recommended for adjusted body weight less than 40 kg(NOTE)*

☐ Antimicrobial - Pharmacy Consult
ONCE
Comments: A pharmacy consult is recommended for adjusted body weight less than 40 kg. Consult to be performed by infectious diseases pharmacist.

CrCl less than 30 mL/min(NOTE)*

☐ DAPTOmycin
300 mg, Soln, IV Push, Every 48 Hour Interval, CrCl less than 30 mL/min
Comments: Adjusted body weight less than 40 kg and creatinine clearance less than 30 mL/min

A pharmacy consult is recommended for adjusted body weight less than 40 kg(NOTE)*

☐ Antimicrobial - Pharmacy Consult
ONCE
Comments: A pharmacy consult is recommended for adjusted body weight less than 40 kg. Consult to be performed by infectious diseases pharmacist.

Adjusted body weight 40 up to 54 kg(NOTE)*
CrCl 30 mL/min or greater(NOTE)*

☐ DAPTOmycin
300 mg, Soln, IV Push, Every 24 Hour Interval, CrCl 30 mL/min or greater
Comments: Adjusted body weight 40 up to 54 kg and creatinine clearance 30 mL/min or greater

CrCl less than 30 mL/min(NOTE)*

☐ DAPTOmycin
300 mg, Soln, IV Push, Every 48 Hour Interval, CrCl less than 30 mL/min
Comments: Adjusted body weight 40 up to 54 kg and creatinine clearance less than 30 mL/min

Adjusted body weight 55 up to 94 kg(NOTE)*
CrCl 30 mL/min or greater(NOTE)*

☐ DAPTOmycin
500 mg, Soln, IV Push, Every 24 Hour Interval, CrCl 30 mL/min or greater
Comments: Adjusted body weight 55 up to 94 kg and creatinine clearance 30 mL/min or greater

CrCl less than 30 mL/min(NOTE)*

☐ DAPTOmycin

500 mg, Soln, IV Push, Every 48 Hour Interval, CrCl less than 30 mL/min

Comments: Adjusted body weight 55 up to 94 kg and creatinine clearance less than 30 mL/min

Adjusted body weight 95 up to 100 kg(NOTE)*

CrCl 30 mL/min or greater(NOTE)*

☐ DAPTOmycin

750 mg, Soln, IV Push, Every 24 Hour Interval, CrCl 30 mL/min or greater

Comments: Adjusted body weight 95 up to 100 kg and creatinine clearance 30 mL/min or greater

CrCl less than 30 mL/min(NOTE)*

☐ DAPTOmycin

750 mg, Soln, IV Push, Every 48 Hour Interval, CrCl less than 30 mL/min

Comments: Adjusted body weight 95 up to 100 kg and creatinine clearance less than 30 mL/min

Adjusted body weight 101 kg or greater(NOTE)*

CrCl 30 mL/min or greater(NOTE)*

☐ DAPTOmycin

750 mg, Soln, IV Push, Every 24 Hour Interval, CrCl 30 mL/min or greater

Comments: Adjusted body weight 101 kg or greater and creatinine clearance 30 mL/min or greater

A pharmacy consult is recommended for adjusted body weight 101 kg or greater(NOTE)*

☐ Antimicrobial - Pharmacy Consult

ONCE

Comments: A pharmacy consult is recommended for adjusted body weight 101 kg or greater. Consult to be performed by infectious diseases pharmacist.

CrCl less than 30 mL/min(NOTE)*

☐ DAPTOmycin

750 mg, Soln, IV Push, Every 48 Hour Interval, CrCl less than 30 mL/min

Comments: Adjusted body weight 101 kg or greater and creatinine clearance less than 30 mL/min

A pharmacy consult is recommended for adjusted body weight 101 kg or greater(NOTE)*

☐ Antimicrobial - Pharmacy Consult

ONCE

Comments: A pharmacy consult is recommended for adjusted body weight 101 kg or greater. Consult to be performed by infectious diseases pharmacist.

Invasive Enterococcus faecium VRE Infections(NOTE)*

Adjusted body weight less than 40 kg(NOTE)*

CrCl 30 mL/min or greater(NOTE)*

☐ DAPTOmycin

500 mg, Soln, IV Push, Every 24 Hour Interval, CrCl 30 mL/min or greater

Comments: Adjusted body weight less than 40 kg and creatinine clearance 30 mL/min or greater

A pharmacy consult is recommended for adjusted body weight less than 40 kg(NOTE)*

☐ Antimicrobial - Pharmacy Consult

ONCE

Comments: A pharmacy consult is recommended for adjusted body weight less than 40 kg. Consult to be performed by infectious diseases pharmacist.

CrCl less than 30 mL/min(NOTE)*

☐ DAPTOmycin

500 mg, Soln, IV Push, Every 48 Hour Interval, CrCl less than 30 mL/min

Comments: Adjusted body weight less than 40 kg and creatinine clearance less than 30 mL/min

A pharmacy consult is recommended for adjusted body weight less than 40 kg(NOTE)*

☐ Antimicrobial - Pharmacy Consult

ONCE

Comments: A pharmacy consult is recommended for adjusted body weight less than 40 kg. Consult to be performed by infectious diseases pharmacist.

Adjusted body weight 40 up to 64 kg(NOTE)*

CrCl 30 mL/min or greater(NOTE)*

☐ DAPTOmycin

500 mg, Soln, IV Push, Every 24 Hour Interval, CrCl 30 mL/min or greater

Comments: Adjusted body weight 40 up to 64 kg and creatinine clearance 30 mL/min or greater

CrCl less than 30 mL/min(NOTE)*

☐ DAPTOmycin

500 mg, Soln, IV Push, Every 48 Hour Interval, CrCl less than 30 mL/min

Comments: Adjusted body weight 40 up to 64 kg and creatinine clearance less than 30 mL/min

Adjusted body weight 65 up to 84 kg(NOTE)*

CrCl 30 mL/min or greater(NOTE)*

☐ DAPTOmycin

750 mg, Soln, IV Push, Every 24 Hour Interval, CrCl 30 mL/min or greater

Comments: Adjusted body weight 65 up to 84 kg and creatinine clearance 30 mL/min or greater

CrCl less than 30 mL/min(NOTE)*

☐ DAPTOmycin

750 mg, Soln, IV Push, Every 48 Hour Interval, CrCl less than 30 mL/min

Comments: Adjusted body weight 65 up to 84 kg and creatinine clearance less than 30 mL/min

Adjusted body weight 85 up to 100 kg(NOTE)*

CrCl 30 mL/min or greater(NOTE)*

☐ DAPTOmycin

1,000 mg, Soln, IV Push, Every 24 Hour Interval, CrCl 30 mL/min or greater

Comments: Adjusted body weight 85 up to 100 kg and creatinine clearance 30 mL/min or greater

CrCl less than 30 mL/min(NOTE)*

☐ DAPTOmycin

1,000 mg, Soln, IV Push, Every 48 Hour Interval, CrCl less than 30 mL/min

Comments: Adjusted body weight 85 up to 100 kg and creatinine clearance less than 30 mL/min

Adjusted body weight 101 kg or greater(NOTE)*

CrCl 30 mL/min or greater(NOTE)*

☐ DAPTOmycin

1,000 mg, Soln, IV Push, Every 24 Hour Interval, CrCl 30 mL/min or greater

Comments: Adjusted body weight 101 kg or greater and creatinine clearance 30 mL/min or greater

A pharmacy consult is recommended for adjusted body weight 101 kg or greater(NOTE)*

☐ Antimicrobial - Pharmacy Consult

ONCE

Comments: A pharmacy consult is recommended for adjusted body weight 101 kg or greater. Consult to be performed by infectious diseases pharmacist.

CrCl less than 30 mL/min(NOTE)*

☐ DAPTOmycin

1,000 mg, Soln, IV Push, Every 48 Hour Interval, CrCl less than 30 mL/min

Comments: Adjusted body weight 101 kg or greater and creatinine clearance less than 30 mL/min

A pharmacy consult is recommended for adjusted body weight 101 kg or greater(NOTE)*

☐ Antimicrobial - Pharmacy Consult

ONCE

Comments: A pharmacy consult is recommended for adjusted body weight 101 kg or greater. Consult to be performed by infectious diseases pharmacist.

Non Categorized

Providers must exercise independent clinical judgment when using order sets.(NOTE)*

***Report Legend:**

DEF - This order sentence is the default for the selected order
GOAL - This component is a goal
IND - This component is an indicator
INT - This component is an intervention
IVS - This component is an IV Set
NOTE - This component is a note
Rx - This component is a prescription
SUB - This component is a subphase