Unique Plan Description: JPP Acute Rehab
Plan Selection Display: JPP Acute Rehab
PlanType: Medical
Version: 18
Begin Effective Date:
End Effective Date: Current
Available at: ***TAMC - THE AROO***
Plan Comment: PRJ0480970_Removed EMMC virtual viewing, added NOTE under Constipation NOTE, added Colace and updated Senokot, lactulose glycerin adult, rectal suppository order comments. Removed Senokot, Fleet Mineral Oil Enema (EMMC), Colace, Provider Communication, Artificial Saliva, lidocaine 4% topical cream, hydrocortisone 25 mg rectal suppository, Reference EMMC (EMMC Only), Vital Signs, Bleach 1:4 diluted cleaning solution (SVH only) orders. Removed Wound Care, Nutrition Therapy, Diagnostic Testing, Respiratory. Miscellaneous., and Therapies sections_RK

JPP Acute Rehab
Patient Care
Medications - Bowel Mgt.

Diarrhea:(NOTE)*

☐ Imodium A-D
2 mg, Cap, PO, Every 3 Hours, PRN, Diarrhea (DEF)*
Comments: Administer after each loose stool; Maximum dose = 16 mg/day, Exclusions: C. diff positive or ruling out C. diff
2 mg, Susp, PO, Every 3 Hours, PRN, Diarrhea
Comments: Administer after each loose stool; Maximum dose = 16 mg/day, Exclusions: C. diff positive or ruling out C. diff

Constipation:(NOTE)*
Start with senna and docusate. If no bowel movement after 24 hours, give lactulose. If still ineffective after another 24 hours, use glycerin suppository.(NOTE)*

☐ Senokot
17.2 mg = 2 TAB, Tab, PO, With Lunch
Comments: For constipation. Start with senna and docusate. If no bowel movement after 24 hours, give lactulose. If still ineffective after another 24 hours, use glycerin suppository. Give with 240 mL of fluid, Exclusions: NPO status

☐ Colace
100 mg, Cap, PO, Daily
Comments: For constipation. Start with senna and docusate. If no bowel movement after 24 hours, give lactulose. If still ineffective after another 24 hours, use glycerin suppository. Give with 240 mL of fluid, For constipation, Exclusions: NPO status

☐ lactulose
20 gm, Syrup, PO, ONCE
Comments: Lactulose 20 gm = 30 mL; For constipation; Start with senna and docusate. If no bowel movement after 24 hours, give lactulose. If still ineffective after another 24 hours, use glycerin suppository.Exclusions: NPO status

☐ glycerin adult rectal suppository
1 SUPP, Supp, RECTAL, Daily, PRN, Constipation
Comments: Start with senna and docusate. If no bowel movement after 24 hours, give lactulose. If still ineffective after another 24 hours, use glycerin suppository. Exclusions: 1. Bowel or prostate surgery within 3 months; 2. Neutropenia (ANC less than 1000); 3. Thrombocytopenia (Platelet count less than 50,000)

Medications - Patient Comfort

Painful hemorrhoids:(NOTE)*

☐ hydrocortisone 2.5% topical cream
1 APP, Cream, TOPICAL, Apply to Perianal, Three Times Daily, PRN, Rectal/Hemorrhoidal Discomfort
Comments: Exclusions: Bowel surgery within past 3 months

Bladder Mgt.

☐ Bladder Scan
PRN Urine Retention
Comments: For Acute urinary retention: Voiding with PVR 150 - 500 mL: Scan in 6 hours. Voiding with PVR greater than 500 mL: Scan in 4 hours. Unable to void, removed 300-500 mL via straight cath: Scan in 6 hours. Unable to void, removed greater than 500 mL via straight

cath: Scan in 4 hours. Discontinue scanning when bladder scan less than 150 mL for 3 consecutive scans and patient voiding more than 180 mL in 5 hours.

☐ Bladder Program (Intermittent Catheterization Program)
Urinary retention

Non Categorized

☑ Subphase (DO NOT COMPLETE)
Providers must exercise independent clinical judgment when using order sets.(NOTE)*

***Report Legend:**
DEF - This order sentence is the default for the selected order
GOAL - This component is a goal
IND - This component is an indicator
INT - This component is an intervention
IVS - This component is an IV Set
NOTE - This component is a note
Rx - This component is a prescription
SUB - This component is a subphase