Unique Plan Description: JPP Med-Surg Adult
Plan Selection Display: JPP Med-Surg Adult
PlanType: Medical
Version: 53
Begin Effective Date:
End Effective Date: Current
Available at: ***BHMH***
C A DEAN HOSPIT
EMMC
MERCY HOSPITAL
NL MAINE COAST HOSPITAL
SVH
TAMC - THE AROO
Plan Comment: PRJ0503294_Removed vitamin E topical oil, nitroglycerin,Alcohol Wipe Sniff,menthol mucous membrane lozenge,Aloe Vesta 2 in 1 Antifungal 2% topical ointment, and heparin flush 100 units/mL intravenous solution orders_RK

JPP Med-Surg Adult
Patient Care

For patients age 18 or greater(NOTE)*

☐ Vital Signs
Every 8 Hours, If patient is afebrile and BP, HR, and RR are within 10% of base line for past 24-hours.
Comments: Exclusions: Critical Care patients

☐ Ocean 0.65% nasal spray
1 Sprays, Soln, NASAL, Four Times Daily, PRN, Dry Nose, 72 hr
Comments: Exclusions: Facial trauma or facial fractures within past 3 months

☐ Artificial Tears ophthalmic solution
1 drops, Soln, BOTH EYES, Every 4 Hours, PRN, Dry Eyes, 72 hr
Comments: Exclusions: Eye Surgery within the last 3 months

☐ Mouth Care
Every 2 hr mouth care with teeth brushing every 12 hr
Comments: Ventilated patient (PRN or nocturnal) with tracheostomy

☐ Heat Application
PRN, PRN Other(comment), Per patient request, Apply on and off in 30 minute intervals
Comments: Exclusions: Confusion/dementia Decreased sensation to requested area.

☐ May Shower
Start: T;N
Comments: Exclusions: Fall Risk (syncope, arrhythmias, deconditioned, oxygen or BiPAP use, tachycardia, tachypnea, hypotensive)

☐ Provider Communication (EMMC)
Patient may be evaluated and treated by certified foot care nurse.

Chest Pain:(NOTE)*

☐ CV ECG
Priority Stat, Reason Chest Pain, ONCE (DEF)*
Priority Stat, Reason Other (Enter in Special Instructions), Telemetry Changes, ONCE

Heartburn:(NOTE)*
Start with calcium carbonate, if ineffective after 2 consecutive doses may give aluminum hydroxide 30 minutes after last calcium carbonate dose(NOTE)*

☐ Tums
500 mg, Chew tab, PO, Every 4 Hours, PRN, Heartburn, 72 hr
Comments: Start with calcium carbonate, if ineffective after 2 consecutive doses may give aluminum hydroxide 30 minutes after last calcium carbonate dose. Exclusions: Diagnosis of renal failure Creatinine Clearance less than 30 mL/min NPO

☐ Mag-al Plus 200 mg-200 mg-20 mg/5 mL oral suspension
30 mL, Susp, PO, Every 4 Hours, PRN, Heartburn, 72 hr
Comments: Start with calcium carbonate, if ineffective after 2 consecutive doses may give aluminum hydroxide 30 minutes after last calcium carbonate dose. Exclusions: Diagnosis of renal failure Creatinine Clearance less than 30 mL/min NPO

Nausea and/or Vomiting:(NOTE)*

☐ Sea Band Acupressure Wristband
Daily, PRN, Nausea
Comments: To P6 point on wrist. May be used any time for any age for nausea and vomiting. May be used alone, in combination, or in addition to medication therapy.

☐ Zofran ODT
4 mg, Disintegrating Tablet, PO, ONCE, Indication: Nausea Indication: Emesis(Vomiting)
Comments: If tolerating PO Exclusions: Post-operative for patient's current admission DO NOT attempt to push ODT tablets through foil backing. With dry hands, peel back the foil of 1 blister and remove the tablet. Place tablet on the tongue; it will dissolve in seconds. Once dissolved, the patient may swallow with saliva. Administration with liquid is not necessary. Wash hands after administration.

☐ Zofran
4 mg, Soln, IV Push, ONCE, Indication: Emesis(Vomiting) Indication: Nausea
Comments: If NPO or not tolerating PO Exclusions: Post-operative for patient's current admission

Hard Stools:(NOTE)*

☐ Colace
100 mg, Cap, PO, Twice Daily, Indication: Hard Stools, 72 hr
Comments: For Hard Stools Exclusions: Rectal and/or Bowel surgery within the last 3 months Absent bowel sounds

Constipation:(NOTE)*

☐ MiraLax
17 gm, Powder, PO, Daily, PRN, Constipation, 72 hr
Comments: If tolerating PO Exclusions: Bowel or prostate surgery within the last 3 months

☐ Dulcolax Laxative
10 mg, Supp, RECTAL, Daily, PRN, Constipation, 72 hr
Comments: If NPO or not tolerating PO. Exclusions: Bowel or prostate surgery within the last 3 months Absolute Neutrophil Count of less than 1,000 Platelet count of 50,000 or less

Pain:(NOTE)*

☐ Tylenol
650 mg, Tablet, PO, Every 4 Hours, PRN, Pain-Mild (Pain Score: 1 - 3), 72 hr
Comments: If tolerating PO Exclusions: NPO status; Abnormal liver enzymes (Total Bilirubin greater than 2 mg/dL, Alkaline Phosphatase greater than 130 units/mL, Alanine Aminotransferase greater than 41 units/mL)

If NPO or not tolerating PO/enteral:(NOTE)*

☐ Tylenol
650 mg, Supp, RECTAL, Every 4 Hours, PRN, Pain-Mild (Pain Score: 1 - 3), 72 hr
Comments: If NPO or not tolerating PO/enteral Exclusions: Patient tolerating PO Abnormal liver enzymes (Total Bilirubin greater than 2 mg/dL, Alkaline Phosphatase greater than 130 units/mL, Alanine Aminotransferase greater than 41 units/mL); Absolute Neutrophil Count of less than 1000; Platelet count of less than or equal to 50,000; Bowel surgery within the last 3 months

If tolerating enteral:(NOTE)*

☐ Tylenol
650 mg, Tablet, NG-TUBE, Every 4 Hours, PRN, Pain-Mild (Pain Score: 1 - 3), 72 hr
Comments: If tolerating enteral Exclusions: NPO status; Abnormal liver enzymes (Total Bilirubin greater than 2 mg/dL, Alkaline Phosphatase greater than 130 units/mL, Alanine Aminotransferase greater than 41 units/mL) Crush and administer via NG-TUBE

Sore Throat:(NOTE)*

☐ Sore Throat Lozenge
1 LOZENGE, Loz, PO, Every 2 Hours, PRN, Sore Throat, 72 hr
Comments: Exclusions: Patients who have received chemotherapy within the last 14 days or if NPO benzocaine-menthol lozenge

Fungal/Skin Irritation:(NOTE)*

☐ miconazole 2% topical powder
1 APP, Powder, TOPICAL, Twice Daily, 7 Days
Comments: Apply to affected areas (Do not apply to feet or open areas) Exclusions: Signs of infection, Absolute Neutrophil Count less than 1,000

☐ Calmoseptine 0.44%-20.6% topical ointment
1 APP, Ointment, TOPICAL, Twice Daily

Comments: Apply to affected area until healed. Indication: Skin irritation

☐ nystatin

500,000 Units, Susp, PO, Four Times Daily, 5 Days, Swish and spit (DEF)*

Comments: Concentration: 100,000 Units = 1 mL Exclusions: Absolute Neutrophil Count less than 1000

500,000 Units, Susp, PO, Four Times Daily, 5 Days, Swish and swallow

Comments: Concentration: 100,000 Units = 1 mL Exclusions: Absolute Neutrophil Count less than 1000; NPO status

IV Therapy - Peripheral:(NOTE)*

☐ lidocaine 1% injectable solution

0.1 mL, Soln, INTRADERMAL, ONCE

Comments: For peripheral IV placement Administering 0.1 mL would deliver 1 mg lidocaine

IV Therapy - Central:(NOTE)*

For thrombotic catheter clearance in non-tunneled lines, including: - PICC (single and double lumen) - Triple lumen (distal, medial, and proximal ports) - Swan Ganz (distal, proximal, and auxiliary lumens)(NOTE)*

☐ Cathflo Activase

2 mg, Soln, CATHETER CLEARANCE, Every 30 Minute Interval, PRN, Catheter Clearance, 7 Days

Comments: For thrombotic catheter clearance Dwell time of two hours minimum Dispense in a 10 mL Luer-lock syringe

☐ alteplase for cath clearance

1 mg, Soln, CATHETER CLEARANCE, Every 30 Minute Interval, PRN, Catheter Clearance, 7 Days

Comments: For thrombotic catheter clearance. Dwell time of 30 minutes minimum, up to 2 hours. May instill a second dose if patency not restored after 2 hours.

For thrombotic catheter clearance in an introducer, tunneled catheter, and/or ports. Examples: - All large bore introducers - Hickman catheter and dialysis tunneled catheters - Regular and high flow ports(NOTE)*

☐ Cathflo Activase

2 mg, Soln, CATHETER CLEARANCE, Every 30 Minute Interval, PRN, Catheter Clearance, 7 Days

Comments: For thrombotic catheter clearance Dwell time of two hours minimum Dispense in a 10 mL Luer-lock syringe

☐ alteplase for cath clearance

2 mg, Soln, CATHETER CLEARANCE, Every 30 Minute Interval, PRN, Catheter Clearance, 7 Days

Comments: For thrombotic catheter clearance. Dwell time of 30 minutes minimum, up to 2 hours. May instill a second dose if patency not restored after 2 hours.

Wound Consult:(NOTE)*

☐ INP Ostomy-Wound Consult

Non-surgical wound, i.e. new skin breakdown, Stage 3 or 4 pressure ulcer, fistula, chronic wound.

Comments: Exclusions: Surgical wound, vascular or plastic surgery patient current admission

Pharmacy Consult:(NOTE)*

☐ Pharmacy Consult - Drug Form or Route Adjustment

ONCE, Comment: Patient unable to tolerate current form of medication(s).

Respiratory:(NOTE)*

☐ Incentive Spirometry

Every 1 Hour Interval, While awake

Sleep:(NOTE)*

☐ melatonin

3 mg, Tab, PO, Bedtime, PRN, Sleep

Comments: Exclusions: NPO status; Other sleeping aids, opiates, muscle relaxants; Decompensated COPD; Morbid obesity requiring CPAP

Non Categorized

☑ Provider Communication

Nursing may administer a lesser potent medication or dose at patient request if an order for PRN pain medication(s) and/or if more than one medication or dose is ordered.

Comments: MUST include "per patient preference" as a comment when documenting administration of lesser potent medication. Example: if 2 or more PRN agents are ordered to treat mild, moderate, and severe pain levels.

☑ Subphase (DO NOT COMPLETE)

Providers must exercise independent clinical judgment when using order sets.(NOTE)*

***Report Legend:**

DEF - This order sentence is the default for the selected order
GOAL - This component is a goal
IND - This component is an indicator
INT - This component is an intervention
IVS - This component is an IV Set
NOTE - This component is a note
Rx - This component is a prescription
SUB - This component is a subphase