Unique Plan Description: Low Dose Ketamine Analgesia Post Op Orders
Plan Selection Display: Low Dose Ketamine Analgesia Post Op Orders
PlanType: Medical
Version: 10
Begin Effective Date:
End Effective Date: Current
Available at: ***MERCY HOSPITAL***
NL MAINE COAST HOSPITAL
Plan Comment:

Low Dose Ketamine Analgesia Post Op Orders
Vital Signs

☑ Vital Signs (Frequent)
Every 1 hour x 4, then every 4 hours
Comments: Vital Signs prior to infusion. Hourly monitoring of vital signs for the first four hours after starting infusion, then every 4 hours until Ketamine infusion discontinued.

Activity

☐ Activity
Other, Advance activity as tolerated. (DEF)*
Bedrest

Patient Care

☑ Agitation Sedation Scale RASS
Every 1 Hour Interval, 4 Doses/Times
Comments: Baseline prior to Ketamine infusion

☑ **+4 Hours** Agitation Sedation Scale RASS
Every 4 Hours
Comments: Until Ketamine infusion discontinued

Continuous Solutions

☐ Sodium Chloride 0.9%
Tot Vol: 1,000 mL, Flow Rate: 20 mL/hr, Soln, IV
Post Op Low Dose Ketamine Infusion:(NOTE)*

☐ ketamine 500 mg / 250 mL NS Pre-Made
Tot Vol: 250 mL, mg/kg/hr, Soln, IV
Comments: CONCENTRATION: 2 mg/mL. (Recommended prescribing range: 0.05 - 0.2 mg/kg/hr). Stop infusion and call Provider for Sedation level of difficult to arouse, following simple commands but drifting off again, no eye contact, sluggish response or sedated. Document in mg/hr. If Bolus is ordered, it may be administered from the smart pump and documented in infusion order.

☐ Ketamine 500 mg/ 250 mL NS (cont) (IVS)*
Sodium Chloride 0.9%
Tot Vol: 250 mL, mg/kg/hr, Soln, IV
Comments: CONCENTRATION: 2 mg/mL. (Recommended prescribing range: 0.05 - 0.2 mg/kg/hr). Stop infusion and call Provider for Sedation level of difficult to arouse, following simple commands but drifting off again, no eye contact, sluggish response or sedated. Document in mg/hr. If Bolus is ordered, it may be administered from the smart pump and documented in infusion order.
ketamine
500 mg = 5 mL

☑ Notify If
Stop Ketamine infusion and call Provider for Sedation level of difficult to arouse, following simple commands but drifting off again, no eye contact, sluggish response or sedated.

☑ Notify If
Page Provider on call if patient becomes dysphoric, agitated or confused.

Respiratory

☑ Oxygen
Via: Nasal Cannula, PRN: Low Oxygen Saturation, Maintain saturation greater than 92%, wean as tolerated.

☑ Oxygen.
Via: Nasal Cannula, PRN: Low Oxygen Saturation, Maintain saturation greater than 92%, wean as

tolerated.

☑ Continuous Pulse Oximetry
Start: T;N

☑ Continuous Pulse Oximetry.
Start: T;N

Non Categorized

Providers must exercise independent clinical judgment when using order sets.(NOTE)*

***Report Legend:**
DEF - This order sentence is the default for the selected order
GOAL - This component is a goal
IND - This component is an indicator
INT - This component is an intervention
IVS - This component is an IV Set
NOTE - This component is a note
Rx - This component is a prescription
SUB - This component is a subphase