Unique Plan Description: OBG Postpartum Hemorrhage
Plan Selection Display: OBG Postpartum Hemorrhage
PlanType: Medical
Version: 15
Begin Effective Date:
End Effective Date: Current
Available at: ***EMMC***
MERCY HOSPITAL
NL MAINE COAST HOSPITAL
TAMC - THE AROO
Plan Comment:

Legend		
Green Highlight	=	Addition
Red Highlight	=	Deletion

OBG Postpartum Hemorrhage
Patient Care

☑ Notify If
Bleeding more than a maternity pad an hour or bleeding appears excessive by clinical judgment

☑ Provider Communication
Place orders for STAT CBC with differential for excessive bleeding and type and screen, PRN if not already obtained.
For excessive bleeding:(NOTE)*

☑ Saline Lock
Ensure patient has 2 patent, large bore (18 g) IVs. One may be saline locked.

Continuous Solutions

☐ Lactated Ringers (Bolus Infusion)
1,000 mL, Soln, IVPB, Post-Procedure, 2 Doses/Times, Rate: 1,000 mL/hr

☐ Sodium Chloride 0.9% (Bolus Infusion)
1,000 mL, Soln, IVPB, Post-Procedure, 2 Doses/Times, Rate: 1,000 mL/hr

Medications

☑ Pitocin
30 Units, Soln, IVPB, ONCE, PRN, Postpartum Bleeding, Infuse Over: 30 Minutes
Comments: LOW RISK Hazardous Drugs At the direction of Provider for Postpartum Hemorrhage
Following delivery of the placenta, for routine management of uterine atony, hemorrhage.(NOTE)*

☑ Cytotec
1,000 mcg, Tab, RECTAL, ONCE, PRN, Bleeding, Indication: Postpartum Uterine Bleeding, Postpartum Hemorrhage
Comments: LOW RISK Hazardous Drugs At the direction of Provider for Postpartum Hemorrhage

☑ Pitocin
10 Units, Soln, IM, ONCE, PRN, Bleeding, Indication: Postpartum Uterine Bleeding, 1 Doses/Times
Comments: LOW RISK Hazardous Drugs At the direction of Provider for Postpartum Hemorrhage

☑ Methergine
0.2 mg, Soln, IM, ONCE, PRN, Bleeding, Indication: Postpartum Uterine Bleeding, 1 Doses/Times
Comments: LOW RISK Hazardous Drugs At the direction of Provider for Postpartum Hemorrhage; Contraindicated in patients with hypertension, pre-eclampsia, pregnancy, and hypersensitivity to methylergonovine (Methergine)

☑ Hemabate
250 mcg, Soln, IM, Every 15 Minute Interval, PRN, Bleeding, Indication: Postpartum Uterine Bleeding, Refractory, 8 Doses/Times
Comments: At the direction of Provider for Postpartum Hemorrhage; Max 8 Doses/Times or 2 mg; Contraindicated in patients with the following: 1. Hypersensitivity to Hemabate (carboprost tromethamine); 2. Acute pelvic inflammatory disease; 3. Patients with active cardiac, pulmonary, renal, or hepatic disease

☑ Imodium A-D
2 mg, Cap, PO, Every 3 Hour Interval, PRN, Diarrhea
Comments: Administer after each loose stool; Max dose = 16 mg/day

☑ tranexamic acid
1 gm, Soln, IVPB, Post-Procedure, PRN, Other(comment), 1 Doses/Times, Infuse Over: 20 10 Minutes

Comments: At the direction of Provider for Postpartum Hemorrhage

Non Categorized

Providers must exercise independent clinical judgment when using order sets.(NOTE)*

***Report Legend:**

DEF - This order sentence is the default for the selected order
GOAL - This component is a goal
IND - This component is an indicator
INT - This component is an intervention
IVS - This component is an IV Set
NOTE - This component is a note
Rx - This component is a prescription
SUB - This component is a subphase