Unique Plan Description: PED JPP
Plan Selection Display: PED JPP
PlanType: Medical
Version: 1
Begin Effective Date:
End Effective Date: Current
Available at: ***EMMC***

PED JPP

Patient Care
Medications - Pain Control

Mild pain or fever greater than 38.5 C:(NOTE)*
Order tablet for patients able to swallow pills, liquid for patients tolerating PO/enteral but unable to swallow pills, suppository for patients unable to tolerate PO/enteral.(NOTE)*
Exclusions: Hepatic insufficiency (Total bilirubin greater than 2 mg/dL, Alanine Aminotransferase greater than 41 units/mL); G6PD Deficiency;(NOTE)*
Dose must be adjusted in patients with renal insufficiency (Serum creatinine greater than 2 mg/dL or Creatinine Clearance less than 30 mL/min). Call provider for dosage(NOTE)*

☐ Tylenol

15 mg/kg, Tablet, PO, Every 6 Hours, PRN, Fever/Pain-Mild (Pain Score: 1 - 3), Weight less than 40 kg [Less Than 40 kg] (DEF)*

Comments: Temperature greater than 38.5 C core or Pain-Mild (Pain Score: 1 - 3); Round dose down to nearest whole tab;
Exclusions: 1. Hepatic insufficiency (Total bilirubin greater than 2 mg/dL, Alanine Aminotransferase greater than 41 units/mL);
2. G6PD Deficiency;
3. Dose must be adjusted in patients with renal insufficiency (Serum creatinine greater than 2 mg/dL or Creatinine Clearance less than 30 mL/min). Call provider for dosage;
4. Do not exceed 3 gm or 75 mg/kg Acetaminophen per 24 hours or 650 mg/dose.
If acetaminophen and ibuprofen ordered together, may alternate.
Weight less than 40 kg

15 mg/kg, Susp, PO, Every 6 Hours, PRN, Fever/Pain-Mild (Pain Score: 1 - 3), Weight less than 40 kg [Less Than 40 kg]

Comments: Temperature greater than 38.5 C core or Pain-Mild (Pain Score: 1 - 3);
Exclusions: 1. Hepatic insufficiency (Total bilirubin greater than 2 mg/dL, Alanine Aminotransferase greater than 41 units/mL);
2. G6PD Deficiency;
3. Dose must be adjusted in patients with renal insufficiency (Serum creatinine greater than 2 mg/dL or Creatinine Clearance less than 30 mL/min). Call provider for dosage;
4. Do not exceed 3 gm or 75 mg/kg Acetaminophen per 24 hours or 650 mg/dose.
If acetaminophen and ibuprofen ordered together, may alternate.
Weight less than 40 kg

15 mg/kg, Supp, RECTAL, Every 6 Hours, PRN, Fever/Pain-Mild (Pain Score: 1 - 3), Weight less than 40 kg [Less Than 40 kg]

Comments: Temperature greater than 38.5 C core or Pain-Mild (Pain Score: 1 - 3); Round dose down to nearest half suppository;
Exclusions: 1. Hepatic insufficiency (Total bilirubin greater than 2 mg/dL, Alanine Aminotransferase greater than 41 units/mL);
2. G6PD Deficiency;
3. Dose must be adjusted in patients with renal insufficiency (Serum creatinine greater than 2 mg/dL or Creatinine Clearance less than 30 mL/min). Call provider for dosage;
4. Do not use suppository in patients with rectal fissure/surgery, neutrophil count less than 1000, or platelet count less than 50,000;
5. Do not exceed 3 gm or 75 mg/kg Acetaminophen per 24 hours or 650 mg/dose.
If acetaminophen and ibuprofen ordered together, may alternate.
Weight less than 40 kg

650 mg, Tablet, PO, Every 6 Hours, PRN, Fever/Pain-Mild (Pain Score: 1 - 3), Weight 40 kg or greater [Greater Than or Equal To 40 kg]

Comments: Temperature greater than 38.5 C core or Pain-Mild (Pain Score: 1 - 3); Round dose down to nearest whole tab;

Exclusions: 1. Hepatic insufficiency (Total bilirubin greater than 2 mg/dL, Alanine Aminotransferase greater than 41 units/mL);
2. G6PD Deficiency;
3. Dose must be adjusted in patients with renal insufficiency (Serum creatinine greater than 2 mg/dL or Creatinine Clearance less than 30 mL/min). Call provider for dosage;
4. Do not exceed 3 gm or 75 mg/kg Acetaminophen per 24 hours or 650 mg/dose. If acetaminophen and ibuprofen ordered together, may alternate.
Weight 40 kg or greater

650 mg, Susp, PO, Every 6 Hours, PRN, Fever/Pain-Mild (Pain Score: 1 - 3), Weight 40 kg or greater [Greater Than or Equal To 40 kg]
Comments: Temperature greater than 38.5 C core or Pain-Mild (Pain Score: 1 - 3);
Exclusions: 1. Hepatic insufficiency (Total bilirubin greater than 2 mg/dL, Alanine Aminotransferase greater than 41 units/mL);
2. G6PD Deficiency;
3. Dose must be adjusted in patients with renal insufficiency (Serum creatinine greater than 2 mg/dL or Creatinine Clearance less than 30 mL/min). Call provider for dosage;
4. Do not exceed 3 gm or 75 mg/kg Acetaminophen per 24 hours or 650 mg/dose. If acetaminophen and ibuprofen ordered together, may alternate.
Weight 40 kg or greater

650 mg, Supp, RECTAL, Every 6 Hours, PRN, Fever/Pain-Mild (Pain Score: 1 - 3), Weight 40 kg or greater [Greater Than or Equal To 40 kg]
Comments: Temperature greater than 38.5 C core or Pain-Mild (Pain Score: 1 - 3); Round dose down to nearest half suppository;
Exclusions: 1. Hepatic insufficiency (Total bilirubin greater than 2 mg/dL, Alanine Aminotransferase greater than 41 units/mL);
2. G6PD Deficiency;
3. Dose must be adjusted in patients with renal insufficiency (Serum creatinine greater than 2 mg/dL or Creatinine Clearance less than 30 mL/min). Call provider for dosage;
4. Do not use suppository in patients with rectal fissure/surgery, neutrophil count less than 1000, or platelet count less than 50,000;
5. Do not exceed 3 gm or 75 mg/kg Acetaminophen per 24 hours or 650 mg/dose. If acetaminophen and ibuprofen ordered together, may alternate.
Weight 40 kg or greater

Medications - Patient Comfort

Heat Application(NOTE)*
For pain or discomfort order either heat application or ice pack based on patient preference(NOTE)*
Exclusions: Patients less than 4 years old or patients unable to provide feedback(NOTE)*

☐ Heat Application
PRN Other(comment), Pain/discomfort
Comments: Apply off and on in 30 minute intervals. If ineffective or patient prefers ice pack, order ice pack.
Exclusions: Patients less than 4 years old or patients unable to provide feedback

Cold Application(NOTE)*
For pain or discomfort order either ice pack or heat application based on patient preference(NOTE)*
Exclusions: Patients less than 4 years old or patients unable to provide feedback(NOTE)*

☐ Ice Pack
PRN Other(comment), Pain/discomfort
Comments: Apply off and on in 30 minute intervals. If ineffective or patient prefers heat application, order heat application.
Exclusions: Patients less than 4 years old or patients unable to provide feedback

Ostomy-Wound Consult for non surgical wound (skin breakdown or stage 3 or 4 pressure ulcer)(NOTE)*

☐ INP Ostomy-Wound Consult

Nasal dryness or infant requiring nasal suctioning of thick secretions:((NOTE)*
Exclusions: Facial trauma or fractures(NOTE)*

☐ Ocean 0.65% nasal spray
1 Sprays, Soln, NASAL, Every 1 Hour Interval, PRN, Dry Nose
Comments: Exclusions: Facial trauma or fractures

Sore throat without signs of infection:(NOTE)*
Exclusions: Patients who have received chemotherapy in the last 7 to 14 days, Patients with risk of aspiration (signs or symptoms of drooling, coughing before or after swallowing, pocketing food inside the cheek, choking on certain foods, trouble chewing or swallowing certain types of fluids or food, gurgling respirations during or after

eating or drinking,(NOTE)*
throat clearing after a meal, frequent coughing when lying flat or sitting up quickly from a reclined position); Patients less than 6 years of age; Abnormal neurologic status (inability to consistently follow verbal commands)(NOTE)*

☐ Sore Throat Lozenge
1 LOZENGE, Loz, PO, Every 2 Hour Interval, PRN, Sore Throat
Comments: Exclusions:
1. Patients who have received chemotherapy in the last 7 to 14 days;
2. Patients with risk of aspiration (Signs or symptoms of drooling, coughing before or after swallowing, pocketing food inside the cheek, choking on certain foods, trouble chewing or swallowing certain types of fluids or food, gurgling respirations during or after eating or drinking, throat clearing after a meal, frequent coughing when lying flat or sitting up quickly from a reclined position);
3. Patients less than 6 years of age;
4. Abnormal neurologic status (inability to consistently follow verbal commands)
benzocaine-menthol lozenge

Difficulty swallowing pills or capsules:(NOTE)*

☐ Pharmacy Consult - Drug Form or Route Adjustment
ONCE, Reason: Other (SEE COMMENT)
Comments: Patient unable to tolerate current form of medication(s)

Venipunctures, port access, or IM injections:(NOTE)*
Exclusions: Child with congenital/idiopathic methemoglobinuria or infants less than 12 months receiving treatment with other methemoglobin inducing agents (sulfa drugs, acetaminophen, nitrofurantoin, nitroglycerin, nitroprusside, phenobarbital, phenytoin);(NOTE)*
G6PD deficiency and in patients with renal/hepatic impairment (Serum creatinine greater than 2 mg/dL, Creatinine Clearance less than 30 mL/min, Total bilirubin greater than 2 mg/dL, Alkaline Phosphatase greater than 130 units/mL, or Alanine Aminotransferase greater than 41 units/mL); Infants less than 1 month of age.(NOTE)*

☐ lidocaine 4% topical cream
1 APP, Cream, TOPICAL, Three Times Daily, PRN, Other(comment) [Greater Than or Equal To 1 Month]
Comments: Apply topically to intact skin 20 minutes prior to venipuncture, port access or IM injection.
Do not swab skin with alcohol prior to applying, Apply cream and massage into site, Cover with occlusive dressing,
Exclusions: 1. Child with congenital/idiopathic methemoglobinuria or infants less than 12 months receiving treatment with other methemoglobin inducing agents (sulfa drugs, acetaminophen, nitrofurantoin, nitroglycerin, nitroprusside, phenobarbital, phenytoin);
2. G6PD deficiency and in patients with renal/hepatic impairment (Serum creatinine greater than 2 mg/dL, Creatinine Clearance less than 30 mL/min, Total bilirubin greater than 2 mg/dL, Alkaline Phosphatase greater than 130 units/mL, or Alanine Aminotransferase greater than 41 units/mL);
3. Infants less than 1 month of age.

Abraded skin:(NOTE)*
Order ophthalmic form for abrasions around eyes and facial abrasions for younger children who may rub eyes(NOTE)*

☐ bacitracin 500 units/g topical ointment
1 APP, Ointment, TOPICAL, Five Times Daily, PRN, Abrasions
Comments: Use ophthalmic form for abrasions around eyes and facial abrasions for younger children who may rub eyes.

☐ AK-Tracin
1 APP, Ointment, TOPICAL, Five Times Daily, PRN, Abrasions
Comments: For abrasions around eyes and facial abrasions for younger children who may rub eyes.

Diaper rash:(NOTE)*
Start with Zinc Oxide, if ineffective order Calmoseptine(NOTE)*

☐ Remedy Zinc Oxide
1 APP, Paste, TOPICAL, Apply to Buttock, Every 1 Hour Interval, PRN, Rash
Comments: Apply to diaper area.
Start with zinc oxide, if ineffective order calmoseptine

☐ Calmoseptine 0.44%-20.6% topical ointment
1 APP, Ointment, TOPICAL, Apply to Buttock, Every 1 Hour Interval, PRN, Rash
Comments: Apply to diaper area with diaper changes

IV Therapy

Need for IV access:(NOTE)*
Exclusion: Central line present(NOTE)*

☐ IV Placement
Insert saline lock for medication access or incompatibility, Exclusions: Central line present

Deaccessing Port-a-cath(NOTE)*

☐ heparin flush 100 units/mL intravenous solution
100 Units, Soln, IV Push, ONCE

Diagnostic Testing

Suspected blood in nasogastric drainage. Notify provider with positive results:(NOTE)*

☐ Gastroccult POC
Start: T;N, ONCE, Expedite

Suspected blood in stools. Notify provider with positive results:(NOTE)*

☐ Hemoccult POC
Start: T;N, ONCE, Expedite

Urine output less than 1 mL/kg/hr or decreased skin turgor/dry mucous membranes. For infants, sunken fontanel:(NOTE)*

☐ Specific Gravity Urine
URINE, Expedite, Start: T;N, ONCE

Altered mental status or reduced oral intake. Notify provider if altered mental status.(NOTE)*

☐ Bedside Glucose Monitoring
ONCE

Respiratory

Decreased oxygen saturations despite interventions (suctioning, position changes, increase HOB)(NOTE)*
Exclusions: Patients with congenital heart disease- provider order for oxygen is required.(NOTE)*

☐ Oxygen.
Via: Nasal Cannula Bubble Humidify, Humidify Oxygen, Keep oxygen saturations greater than 90% using appropriate humidified oxygen device. Notify provider. Exclusions: Patients with congenital heart disease - provider order for oxygen required

Poor respiratory effort(NOTE)*
For patients too young to follow instructions for incentive spirometry, utilize bubbles or windmill instead.(NOTE)*

☐ Incentive Spirometry
10 times per hour

MDI administration:(NOTE)*

☐ Aerochamber Inhaler Device
1 EA, Device, INHALATION, Daily, PRN, Use With Inhaler

GI

Consider non-medication interventions such as Alcohol wipe sniff, Sea band acupressure or Peppermint oil in room, for nausea management. If ineffective, contact provider for additional guidance(NOTE)*
No bowel movement in past 48 hours:(NOTE)*

☐ glycerin pediatric rectal suppository
1 SUPP, Suppository, RECTAL, ONCE [Less Than 6 YR]
Comments: If no bowel movement in past 48 hours, For ages less than 6 years, Exclusions: 1. Immunocompromised patient, 2. Rectal fissures or rectal surgery.
If ineffective, discuss with provider

☐ glycerin adult rectal suppository
1 SUPP, Suppository, RECTAL, ONCE [Greater Than or Equal To 6 YR]
Comments: If no bowel movement in past 48 hours, For age 6 years and greater, Exclusions: 1. Immunocompromised patient, 2. Rectal fissures or rectal surgery. If ineffective, discuss with provider

Non Categorized

☑ Subphase (DO NOT COMPLETE)
Providers must exercise independent clinical judgment when using order sets.(NOTE)*

***Report Legend:**
DEF - This order sentence is the default for the selected order

GOAL - This component is a goal
IND - This component is an indicator
INT - This component is an intervention
IVS - This component is an IV Set
NOTE - This component is a note
Rx - This component is a prescription
SUB - This component is a subphase